

Copper(I) complexes of *N*-thioacylamido(thio)phosphates and triphenylphosphine

Alexander Y. Verat^{a,*}, Felix D. Sokolov^{a,*}, Nail G. Zabiroy^a, Maria G. Babashkina^a,
Dmitry B. Krivolapov^b, Vasilii V. Brusko^a, Igor A. Litvinov^b

^a A.M. Butlerov Chemical Institute, Kazan State University, 18 Kremlevskaya street, Kazan 420008, Russian Federation

^b A.E. Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Centre of the Russian Academy of Sciences, 8 Arbuzov street, Kazan 420088, Russian Federation

Received 21 May 2005; received in revised form 21 September 2005; accepted 22 September 2005

Available online 28 November 2005

Abstract

Heteroligand copper(I) complexes of bi- or bis-bidentate acylamidophosphates $\text{PhC(S)NHP(S)(OPr-i)}_2$, $\text{PhC(S)NHP(O)(OPr-i)}_2$, $\text{Et}_2\text{NC(S)NHP(S)(OPr-i)}_2$, $\text{PhNHC(S)NHP(S)(OPr-i)}_2$, *N*-(4-aminobenzo-15-crown-5)- $\text{C(S)NHP(S)(OPr-i)}_2$, *N,N*-(1,10-diaza-18-crown-6)- $[\text{C(S)NHP(S)(OPr-i)}_2]_2$, and triphenylphosphine were prepared and characterised. Copper is bound by two PPh_3 and one SCNPX ($\text{X} = \text{O}, \text{S}$) fragment of chelating ligand in all cases. Triphenylphosphine molecules reversibly dissociate in solution. Details of the X-ray structures of $(\text{Ph}_3\text{P})_2\text{Cu}[\text{PhC(S)NP(S)(OPr-i)}_2]$ and $(\text{Ph}_3\text{P})_2\text{Cu}[\text{Et}_2\text{NC(S)NP(S)(OPr-i)}_2]$ are reported.

© 2005 Elsevier B.V. All rights reserved.


Keywords: Copper complexes; *N*-phosphorylated thioureas; *N*-phosphorylated amides; Triphenylphosphine; Crystal structures

1. Introduction

N-(thio)phosphorylated (thio)amides (**1**) and *N*-(thio)phosphorylated (thio)ureas (**2**) are known to form complexes with a variety of metals both soft and hard. Many complexes of transition [1–5] and alkaline [5–9] metals with *N*-acylamidophosphinates (the general name of compounds containing XCNPY backbone) have been reported (see Scheme 1). Among them, transition metal complexes with C(S)NHP(S) ligands have been investigated most extensively.

Cu(I) and Ag(I) complexes with **1** and **2** are polynuclear in solid state. Thus, Ag(I) cations with thiobenzamide $\text{PhC(S)NHP(S)(OPr-i)}_2$ (**3**) form a tetrameric cyclic complex (**4**) [10]. The silver atom in **4** has trigonal configuration. A similar cyclic structure has been found in the trimeric complex of Cu(I) (**5**), obtained with the *N*-thio-

phosphorylated thiourea **2a** (both $\text{R} = \text{Et}$; $\text{R}' = \text{OPh}$) [11]. The copper atoms of **5** adopt a trigonal configuration as well. The alteration in the ring size for **4** and **5** in comparison with each other can be explained both by the difference in ligand structure and in ionic radii of Cu(I) and Ag(I) cations.



Complexes of monovalent copper are of interest as catalysts of numerous processes including homolytic C–Hal (Hal = Cl, Br) bond cleavage in polyhaloalkanes [12]. *N*-thiophosphorylated bis-thioureas containing a crown ether

* Corresponding authors. Tel.: +7 8432 315310; fax: +7 8435 543754.

E-mail addresses: averat@mail.ru (A.Y. Verat), felix.sokolov@ksu.ru (F.D. Sokolov).